

X-ray and Near-infrared Studies of a Star-forming Cloud; L1448

M. Tsujimoto^{1,2}, N. Kobayashi³, & Y. Tsuboi⁴

ABSTRACT

We present the results of X-ray and near-infrared (NIR) observations of L1448, a star-forming region in the Perseus cloud complex using the *Chandra* X-ray Observatory and the 4 m telescope at the Kitt Peak National Observatory. We detect 72 X-ray sources in a $\sim 17' \times 17'$ region with a ~ 68 ks ACIS exposure, for which we conduct follow-up NIR imaging observations in a concentric $\sim 11' \times 11'$ region with FLAMINGOS down to $m_{K_S} \sim 17$ mag. Twelve X-ray sources have NIR or optical counterparts. By plotting X-ray mean energy versus NIR to X-ray flux ratio, the X-ray sources are clearly separated into two groups. The X-ray spectral and temporal features as well as NIR magnitudes and colors indicate that one group mainly consists of young stellar objects (YSOs) in the cloud and the other of background extragalactic sources. Ten X-ray-emitting YSO candidates are thus newly identified, which are low-mass or brown dwarf mass sources from their NIR magnitudes. In addition, a possible X-ray signal is found from a mid-infrared protostar L1448 IRS 3(A). The lack of detection of this source in our deep NIR images indicates that this source has a very steep spectral slope of $\gtrsim 3.2$ in $2\text{--}10\ \mu\text{m}$.

Subject headings: X-rays: stars — infrared: stars — stars: pre-main-sequence — stars: individual (L1448)

1. INTRODUCTION

L1448 is a dark cloud (Lynds 1962) located at the western edge of the Perseus molecular cloud complex at a distance of ~ 300 pc. It has a size of $\sim 15' \times 7'$ in a C^{18}O map and a mass of $\sim 100\ M_{\odot}$ (Bachiller & Cernicharo 1986). The region has been intensively studied in wavelengths longer than far-infrared focusing mainly on a giant molecular outflow, and only few studies in the near-infrared (NIR) broad-band (Bally et al. 1993) and none in the X-ray band have been presented, both of which are currently strong measures to

identify stellar constituents of star-forming clouds. The infrared excess due to emission from circumstellar material and the X-ray brightness are two major observational characteristics of young stellar objects (YSOs), which can be respectively revealed through NIR broad-band photometry in the J , H , and K_S bands and X-ray imaging observations. The purpose of this paper is to report the results of the first X-ray imaging-spectroscopy study of L1448 aided by follow-up NIR broad-band imaging observations. Previous X-ray studies in the Perseus cloud complex can be found in Preibisch et al. (1996); Preibisch & Zinnecker (2001, 2002, 2004) for IC 348, Getman et al. (2002); Preibisch (2003) for NGC 1333, and Yamauchi et al. (2001) for Barnard 1 and IRAS 03282+3035.

The most magnificent feature in L1448 is a parsec-scale bipolar outflow emanating from the core of the cloud, which is one of the highest-velocity, best-collimated, and youngest outflows known (Davis & Smith 1995). The outflow was discovered by Bachiller et al. (1990) in a molecular emission line image. The exciting source of

¹Visiting Astronomer, Kitt Peak National Observatory, National Optical Astronomy Observatory, which is operated by the Association of Universities for Research in Astronomy, Inc. (AURA) under cooperative agreement with the National Science Foundation.

²Department of Astronomy & Astrophysics, The Pennsylvania State University, 525 Davey Laboratory, University Park, PA 16802.

³Institute of Astronomy, The University of Tokyo, 2-21-1 Osawa, Mitaka, Tokyo 181-0015, Japan.

⁴Department of Physics, Chuo University, Kasuga 1-13-27, Bunkyo-ku, Tokyo 112-8551, Japan.

the outflow was promptly detected in the millimeter and centimeter continua (Bachiller et al. 1991; Curiel et al. 1990), which is referred as L1448 mm or L1448 C(enter). Barsony et al. (1998) obtained a spectral energy distribution (SED) in far-infrared to millimeter wavelengths and established the class 0 status of this source.

The *Infrared Astronomical Satellite* (IRAS) detected three sources (IRAS 03220+3035, 03222+3034, and 03225+3034) in this region, which are called L1448 IRS 1, 2, and 3. IRS 1 has a visual counterpart with $m_V = 19.5$ mag (Cohen & Kuhi 1979). From its strong H α emission and excess emission in the *IRAS* band (Cohen & Kuhi 1979; de Grijp et al. 1987), it is probably a classical T Tauri star or a class I protostar in the cloud. IRS 2 is a class 0 source confirmed by O’Linger et al. (1999) based on an SED using *IRAS* and SCUBA data. This source is associated with various signatures of extreme youth, including CO and H₂ molecular outflows (O’Linger et al. 1999; Eislöffel 2000), Herbig-Haro objects (Bally et al. 1997), an H₂O maser source (Anglada et al. 1989), and a centimeter continuum emission (Anglada & Rodríguez 2002). IRS 3, also known as L1448 N(orth), is resolved into three components; IRS 3(A), 3(B), and 3(C), also known respectively as L1448 N(A), N(B), and NW. The former two constitute a binary separated by ~ 2100 AU. Classification of the individual binary components in millimeter to far-infrared observations had been difficult due to crowding. However, recent mid-infrared (MIR) observations clearly resolved IRS 3(A) and 3(B) with an *N*-band detection only from 3(A). Based on this, Ciardi et al. (2003) argued that 3(B) is a class 0 protostar whereas 3(A) might be at the transition phase between class 0 and I stages. IRS 3(C), away from the binary by $\sim 20''$, is another class 0 source (Barsony et al. 1998).

2. OBSERVATIONS

The X-ray observation was carried out on 2004 March 15 using ACIS (Advanced CCD Imaging Spectrometer; Garmire et al. 2003) onboard *Chandra* (Weisskopf et al. 2002). Four ACIS-I chips (I0, I1, I2, and I3) covered a $\sim 17' \times 17'$ region aimed at R. A. = $3^h 25^m 36.4^s$ and Decl. = $30^\circ 44' 58''$ (J2000.0) for an exposure time of 67.9 ks. ACIS has sensitivity in the 0.5–9.0 keV

band with a resolution of ~ 150 eV at 6 keV and a superb point spread function (PSF) radius of $\sim 0.5''$ at the on-axis position. The data were taken with the very faint telemetry mode and the timed exposure CCD operation with a frame time of 3.2 s.

The NIR observation was conducted on 2003 December 16 using FLAMINGOS (Elston et al. 2003) on the Cassegrain focus of the 4 m telescope at the Kitt Peak National Observatory. FLAMINGOS uses a HgCdTe HAWAII-2 array which has a format of 2048×2048 with a pixel scale of $\sim 0.316''$ pixel⁻¹. With dithering observations of an amplitude of $15''$, we obtained *J*-, *H*-, and *K_S*-band images that cover a $\sim 11' \times 11'$ area centered close to the ACIS aim point at R. A. = $3^h 25^m 35.1^s$ and Decl. = $30^\circ 45' 19''$ (J2000.0). The total exposure time was 13.5, 7, and 17 minutes respectively in the *J*, *H*, and *K_S* bands. The seeing was $\sim 1''$. Figures 1 (a) and (b) show the X-ray and NIR intensity maps. The outflow from L1448 mm appears clearly in the *K_S* band (Fig. 1b) presumably due to H₂ $v=1-0$ S(1), vibrational-rotational transition line emission by excited hydrogen molecules.

3. DATA REDUCTION & ANALYSIS

3.1. X-ray Data

We reprocessed the Level 1 data distributed by the *Chandra* X-ray Center to obtain our X-ray event list. The latest calibration results (CALDB 2.28) were incorporated and the background rejection algorithm specific to data taken with the very faint mode was applied. Events were further cleaned by removing cosmic-ray afterglows and applying filters based on the event grade, status, and good-time intervals.

We detected sources using a wavelet technique in CIAO¹ from three images of different energy bands; soft (0.5–2.0 keV), hard (2.0–9.0 keV), and total (0.5–9.0 keV). For each detected source, we used ACIS Extract² for a systematic event extraction, background subtraction, calculation of instrumental responses, and construction and binning of spectra and light curves. Source events were extracted from a 90% encircled energy poly-

¹See <http://asc.harvard.edu/ciao/> for detail.

²See http://www.astro.psu.edu/xray/docs/TARA/ae_users_guide.html for detail.




Fig. 1.— (a) X-ray image. The 0.5–9.0 keV intensity is given by a logarithmic grey scale. The positions of X-ray sources with and without NIR counterparts are shown by crosses and pluses respectively, whereas the position of three IRAS sources (IRS 1–3) and L1448 mm are by the filled squares. The contours indicate the intensity of an NH₃ transition line (Anglada et al. 1989). The fields of view of four ACIS-I chips and FLAMINGOS are illustrated with the solid oblique squares and the dotted square. (b) NIR image. The K_S-band intensity is given by a logarithmic grey scale. The positions of embedded sources are bracketed with a pair of solid lines when undetected and by crosses when detected.

gon of the 1.5 keV PSF, whereas background events were locally extracted from an annular region around the source. The positions of sources were fine-tuned by correlating the event distribution with the PSF. As the X-ray source distribution is relatively sparse in the study field (Fig. 1a), we do not suffer any overlaps of extraction regions. The details of the procedure are described in Getman et al. (2005).

For each source, we derived X-ray photometry information consisting of the source position and its uncertainty, source count (0.5–8.0 keV), net count rate (NCR), probability of no source (PNS), photometry significance (PS), and mean energy (ME). The net count (C_{net}) is calculated from the number of counts (C_{src} and C_{bkg}) and the extraction area (A_{src} and A_{bkg}) of the source and background regions as $C_{\text{net}} = C_{\text{src}} - C_{\text{bkg}} \times (A_{\text{src}}/A_{\text{bkg}})$, which is divided by the effective exposure time to derive NCR. PNS is an index for the detection significance, with which we test the null hypothesis that all detected counts are explained by background fluctuation;

$$\text{PNS} = 1 - \int_0^{C_{\text{src}}-1} dN P(C_{\text{bkg}} \left(\frac{A_{\text{src}}}{A_{\text{bkg}}} \right), N), \quad (1)$$

where $P(\lambda, N)$ represents a Poisson probability distribution function of a mean λ to have N counts. PS is a metric for the reliability of photometry defined as

$$\text{PS} = \frac{C_{\text{net}}}{\sqrt{(\Delta C_{\text{src}})^2 + (\Delta C_{\text{bkg}})^2 \times (A_{\text{src}}/A_{\text{bkg}})^2}}, \quad (2)$$

where the approximation by Gehrels (1986) is used to estimate $\Delta C_{\text{src}} = 1 + \sqrt{C_{\text{src}} + 0.75}$ and $\Delta C_{\text{bkg}} = 1 + \sqrt{C_{\text{bkg}} + 0.75}$. ME is a metric to represent the spectral hardness of sources. The value is the average incident energy (E) weighted by C_{net} of each energy bin and is defined as

$$\text{ME} = \frac{\int dE E C_{\text{net}}(E)}{\int dE C_{\text{net}}(E)}. \quad (3)$$

Table 1 shows the results of the ACIS photometry, consisting of 72 sources with high significance both in the detection ($\text{PNS} \leq 1 \times 10^{-3}$) and the photometry ($\text{PS} \geq 2$).

For each source, flux variability was tested using the K-S test and the Bayesian block segmentation technique (Scargle 1998). We recognized flux

TABLE 1
ACIS PHOTOMETRY

X-ray ^a ID	R.A. ^b (hh:mm:ss.s)	Decl. ^b (dd:mm:ss)	Error ($''$)	Counts	NCR ^c (ks ⁻¹)	PNS ^c	PS ^c	ME ^c (keV)	NIR ^d Counterpart
1	3:24:59.3	30:44:41	0.7	25	0.3	0	3.2	3.6	...
2	3:25:02.7	30:38:34	1.1	25	0.6	9e-7	2.8	3.3	...
3	3:25:03.8	30:43:28	0.6	20	0.3	0	3.0	2.7	...
4 [†]	3:25:04.8	30:38:58	0.4	116	2.4	0	9.0	1.4	03250474+3038578
5	3:25:07.2	30:38:04	1.0	34	0.6	0	3.8	3.6	...

^aVariable sources are marked with a dagger.

^bEquinox in J2000.0.

^cSee text for definitions of NCR, PNS, PS, and ME. PNS values are replaces with 0 when $\text{PNS} < 1\text{e}-9$.

^d2MASS counterparts are given in the format of hhmmssss+ddmmss, while FLAMINGOS counterparts are given by their source numbers in Table 3.

NOTE.—The complete version of this table is in the electronic edition of the Journal. The printed edition contains only a sample.

to be variable if the number of Bayesian blocks exceeds two and the K-S probability is less than 1%. Four sources (Nos. 4, 7, 34, and 69) have variable flux, which are marked with daggers in Table 1.

For 18 sources with more than 40 counts, we performed spectral analysis. The spectra were fit with a one-temperature thin-thermal plasma model (the APEC model; Smith et al. 2001) with the metallicity value fixed to 0.3 solar to determine the best-fit absorption (N_{H}), plasma temperature ($k_{\text{B}}T$), and X-ray luminosity (L_{X}) values. They were also fit with a power-law model to derive the best-fit N_{H} , index (Γ), and absorption-corrected flux ($F_{\text{X}}^{\text{cor}}$) values. Table 2 shows the results of 13 and 11 successful fits with the thermal and power-law models with a reasonable value of $k_{\text{B}}T < 10$ keV and $\Gamma < 3$.

3.2. NIR Data

All FLAMINGOS frames were reduced following the standard procedures using IRAF; i.e., subtraction of dark current, flat-fielding using a dome flat, subtraction of sky using median sky, removal of bad pixels, and trimming of edges. Several frames with readout failure were discarded. Artificial signals were masked out, which we found at pixels to the right and left of, or above and below bright sources by ~ 128 pixels. Nebulosity extending larger than the dithering amplitude may remain in the median-sky-subtracted images; it does not affect our photometry. Referring to sources in

the 2MASS all-sky survey catalog (Skrutskie et al. 1997), all FLAMINGOS frames were corrected for astrometry by a low-order polynomial function to compensate for the image distortion. They were also corrected for photometry to match 2MASS magnitudes. After these corrections, the frames were combined into the final J -, H -, K_{S} -band images.

We extracted sources from the K_{S} -band image (Fig. 1b) above a 5σ level using SExtractor (Bertin & Arnouts 1996). The number of false positive detections is statistically negligible. After manually removing suspicious sources (introduced mostly by misidentifying outflow knots as point sources) and sources at the edges, we obtained 294 detections. We measured J -, H -, and K_{S} -band magnitudes at their positions (Table 3). We matched the FLAMINGOS and the 2MASS sources and found that all the 67 2MASS sources in the FLAMINGOS field of view have FLAMINGOS counterparts.

Figure 2 shows the astrometric accuracy of the FLAMINGOS sources, where the difference of FLAMINGOS and 2MASS positions ($\Delta\text{R.A.}$ and $\Delta\text{Decl.}$) are plotted for the 67 FLAMINGOS–2MASS counterpart pairs. The root mean square of $\Delta\text{R.A.}$ and $\Delta\text{Decl.}$ are $0.18''$ and $0.33''$, indicating that the FLAMINGOS position accuracy is $\sim 0.4''$ (≈ 1.2 FLAMINGOS pixel). The accuracy is better for most sources, as the values are exaggerated by three sources at the top left in Fig-

TABLE 2
ACIS SPECTROSCOPY OF BRIGHT SOURCES

X-ray ID	thermal model			power-law model		
	$N_{\mathrm{H}}^{\mathrm{a}}$ (10^{22} cm^{-2})	$k_{\mathrm{B}}T^{\mathrm{a}}$ (keV)	$L_{\mathrm{X}}^{\mathrm{a},\mathrm{b}}$ ($10^{27} \text{ ergs s}^{-1}$)	$N_{\mathrm{H}}^{\mathrm{a}}$ (10^{22} cm^{-2})	Γ^{a}	$F_{\mathrm{X}}^{\mathrm{cor a},\mathrm{b}}$ ($10^{-14} \text{ ergs s}^{-1} \text{ cm}^{-2}$)
4	0.0 (0.0–0.1)	1.7 (1.5–1.9)	1.8 (1.1–2.0)	...	...	...
6	0.4 (0.2–0.6)	0.9 (0.7–1.1)	1.3 (0.0–1.6)	...	...	...
10	1.3 (0.8–2.0)	2.7 (1.4–5.9)	3.6 (0.2–3.8)	1.6 (1.3–2.3)	2.7 (2.3–3.3)	5.7 (0.0–6.7)
11	...	...	...	0.3 (0.2–0.6)	1.2 (0.9–1.5)	4.1 (0.0–4.6)
15	0.0 (0.0–0.2)	1.0 (0.9–1.2)	1.0 (0.0–1.5)	...	...	...
16	0.3 (0.2–0.4)	5.3 (4.3–7.1)	24.9 (21.7–26.5)	0.4 (0.3–0.5)	1.9 (1.8–2.0)	26.1 (21.7–29.0)
20	0.3 (0.2–0.3)	7.9 (5.5–11.2)	10.2 (8.3–10.7)	0.3 (0.3–0.4)	1.7 (1.6–1.8)	10.1 (7.8–11.4)
23	0.3 (0.1–0.5)	0.6 (0.5–0.8)	0.9 (0.4–1.3)	...	...	...
42	0.0 (0.0–0.1)	0.6 (0.5–0.6)	2.1 (0.8–2.0)	...	...	...
48	0.9 (0.7–1.0)	0.6 (0.5–0.6)	7.7 (3.6–12.4)	...	...	...
50	...	...	...	0.1 (0.0–0.3)	1.4 (1.0–1.6)	1.3 (0.0–1.5)
53	...	...	...	0.4 (0.1–0.9)	1.1 (0.6–1.6)	1.4 (0.0–1.6)
54	...	...	...	0.5 (0.3–1.0)	1.6 (1.2–2.1)	1.7 (0.0–2.1)
64	0.8 (0.6–1.0)	5.7 (3.8–9.9)	7.1 (3.8–7.4)	0.9 (0.7–1.2)	1.9 (1.7–2.1)	7.5 (3.6–8.8)
66	0.4 (0.2–0.7)	3.2 (2.0–6.4)	1.3 (0.3–1.7)	0.5 (0.3–0.9)	2.3 (1.8–2.9)	1.5 (0.0–1.9)
68	2.1 (1.4–3.1)	2.0 (1.2–4.7)	3.9 (0.2–4.3)	...	...	...
71	0.8 (0.3–1.4)	5.7 (2.3–15.0)	3.3 (0.0–3.7)	0.8 (0.5–1.6)	1.8 (1.3–2.7)	3.3 (0.0–4.3)
72	...	...	...	1.0 (0.8–1.4)	1.6 (1.3–1.9)	5.1 (0.0–5.9)

^aThe 1 σ confidence ranges are given in the parentheses.

^bAbsorption-corrected values in the 0.5–8.0 keV band. A distance of 300 pc is assumed to derive L_{X} for the thermal fits.

TABLE 3
FLAMINGOS PHOTOMETRY

NIR ^a ID	R.A. ^b (hh:mm:ss.s)	Decl. ^b (dd:mm:ss)	J^{c} (mag)	H^{c} (mag)	K_{S} (mag)	2MASS Counterpart
1	3:25:09.4	30:48:59	18.8	18.1	17.1	...
2	3:25:09.5	30:46:21	12.5	11.0	10.1	03250943+3046215
3	3:25:09.7	30:48:48	13.3	12.5	12.2	03250965+3048488
4	3:25:10.1	30:46:07	16.4	16.2	17.6	...
5	3:25:10.2	30:47:13	...	...	16.5	...

^aSources with NIR excess are marked with a dagger.

^bEquinox in J2000.0.

^cMagnitudes for sources detected with more than 3 σ are given.

NOTE.—The complete version of this table is in the electronic edition of the Journal. The printed edition contains only a sample.

ure 2, all of which are located too far away from the image center to be corrected for distortion.




Fig. 2.— Astrometric accuracy of FLAMINGOS sources. The difference of R.A. and Decl. between the FLAMINGOS and 2MASS positions (Δ R.A. and Δ Decl.) are plotted for all the FLAMINGOS–2MASS pairs. The solid square at the center represents the FLAMINGOS pixel size of $0.316'' \times 0.316''$.

We estimated the detection completeness of the FLAMINGOS images by embedding artificial sources and detecting them with the same source detection algorithm. The detection rates at different magnitude bins were derived to estimate the 90% completeness limit as ~ 18.0 mag, ~ 16.5 mag, and ~ 17.0 mag in the J , H , and K_S bands.

3.3. NIR and Optical Identification of X-ray Sources

Seven X-ray sources were found to have FLAMINGOS counterparts within $1.5''$ and five additional sources to have 2MASS counterparts outside of the FLAMINGOS view. X-ray positions in Table 1 are shifted by $\sim 0.3''$ to match the 2MASS astrometry. The root mean square of the positional difference between 2MASS–X-ray pairs is $\sim 0.6''$ (≈ 1.3 ACIS pixel). Ten of the NIR-identified X-ray sources are optically identified with the USNO-B catalog (Monet et al. 2003). None of the NIR-unidentified

X-ray sources were identified in the optical catalog. No X-ray emission was found from Herbig-Haro objects (Bally et al. 1997) or H_2 knots (Davis & Smith 1995).

4. DISCUSSION

4.1. Nature of X-ray Sources

Two independent quantities from the X-ray photometry (ME and NCR) combined with the NIR identification give a clue to classify and reveal the nature of the 72 X-ray sources. Figure 3 shows a scatter plot of ME and NCR. Most of the X-ray sources with NIR counterparts (*filled symbols*) are distributed in the limited ME range of $\lesssim 2.5$ keV, while those without NIR counterparts (*open symbols*) are distributed in the limited ME and NCR ranges of $\gtrsim 2.0$ keV and $\lesssim 1.0$ ks^{-1} . This implies that the X-ray sources consist of two major classes of different nature.




Fig. 3.— Scatter plot of ME and NCR for all X-ray sources. Sources with and without NIR counterparts are shown with filled and open symbols, respectively. The source numbers (Table 1) of the former are labeled in Roman, while the latter is in *italic* if they have ≥ 40 counts. The median value of ME and NCR are shown with dashed-and-dotted lines.

The separation of the two classes becomes more apparent when NIR flux is taken into account.

Figure 4 shows a scatter plot of ME and the ratio of NIR flux in the K_S band to X-ray flux in 0.5–8.0 keV band. Here, the X-ray flux (F_X) is calculated by

$$F_X = \frac{\int dE E C_{\text{net}}(E)}{EA t_{\text{exp}}} = \frac{\text{ME} \times \text{NCR}}{EA}, \quad (4)$$

where t_{exp} is the exposure time and EA is the effective area of the mirror averaged over the energy. The X-ray sources with NIR counterparts (*large filled symbols*) and those without FLAMINGOS counterparts (*large open squares*) are clearly separated into two groups. One group (group A) is comprised of X-ray source Nos. 4, 6, 7, 14, 15, 23, 30, 42, and 48 in the upper left quarter, while the other (group B) includes Nos. 11, 16, 39, and FLAMINGOS-unidentified X-ray sources (*large open squares*) at the bottom. NIR-unidentified X-ray sources outside of the FLAMINGOS field of view (*small open symbols*) are hard to classify into either group as the K_S -band upper limit of 2MASS is not deep enough. This illustrates the importance of our NIR observation deeper than 2MASS, even if it results in no detection from X-ray sources.

We conclude that the sources in group A are mostly YSOs that belong to L1448, while those in group B are background AGNs. We note that NIR identification of an X-ray source in star-forming regions does not always guarantee its stellar origin because AGNs, which constitutes the largest contaminant, can be detected even at the 2MASS depth. Instead, we employ the NIR to X-ray flux ratio as an effective measure to separate these two classes. The optical to soft X-ray flux ratio was used for a similar classification purpose in X-ray blank-sky surveys by *Einstein* and *ROSAT* satellites (Maccacaro et al. 1988; Krautter et al. 1999). We use the NIR to total band X-ray flux to be suitable for star-forming regions, which are routinely highly obscured. In (low-mass) main sequence and pre-main-sequence (PMS) sources, X-ray emission originates from stellar coronae while NIR emission originates from photosphere and from circumstellar matter. In AGNs, on the other hand, X-ray emission is from accretion disks around the central black hole while NIR is from hot dust and jets. The different origins bring different typical values of NIR to X-ray flux ratio in stars and AGNs.

Several lines of evidence support this conclu-




Fig. 4.— Scatter plot of ME and the flux ratio of NIR (K_S) and X-ray (0.5–8.0 keV) bands for all X-ray sources with and without NIR counterparts (*filled* and *open*). Open symbols have either a normal size (for sources inside the FLAMINGOS field of view) or a smaller size (otherwise). Variable sources are marked with diamonds. Open symbols indicate the upper limit arising from no K_S -band detection. Labels are given in the same manner with Fig. 3. Variable sources are also labeled. The median values of ME is shown with the dashed-and-dotted line.

sion. First, all sources with ≥ 40 counts in group A (Nos. 4, 6, 15, 23, 42, and 48) have a successful fit with thermal spectral models (Table 2) but not with power-law models with reasonable parameters. On the other hand, all sources with ≥ 40 counts in group B (Nos. 11, 16, 50, and 53) have a successful fit with power-law models. Three (Nos. 11, 50, and 53) of them are not fit with thermal models (Table 2). The clear distinction in X-ray spectral features is consistent with the idea that (1) these groups of sources are different in nature, (2) group A sources are stellar X-ray emission, which typically show thermal spectra of $k_B T \sim 1$ keV, and (3) group B sources are AGNs, which typically show power-law spectra of $\Gamma \sim 1.7$. This conversely enables us to use X-ray spectral features to classify sources for which it is unclear which group to belong to. Nos. 54 and 72 are group B sources for successful fits only by the power-law model (Table 2).

The second argument is the X-ray flux variability of a typical time scale of ~ 10 ks. Stellar X-ray emission from coronae is often characterized by X-ray flares due to magnetic reconnection, whereas no such short time-scale variability is expected from AGNs. Sources with X-ray variability are seen in group A (Nos. 4 and 7) and not in group B. This conversely indicates that sources with similar variability (Nos. 34 and 69) belong to group A.

Third, NIR magnitudes of the X-ray sources in group A are distributed in $m_{K_S} \sim 9\text{--}13$ mag. Contamination by extragalactic sources in this magnitude range is negligible as the expected number of galaxies in the FLAMINGOS field of view is $\lesssim 1$ in $m_{K_S} \lesssim 14$ mag (Tokunaga 2000). The color-magnitude diagram (Fig. 5) shows that group A sources have consistent NIR magnitudes for low mass or very low mass stars at a distance of 300 pc. Sources in group B, on the contrary, have $m_{K_S} \gtrsim 15$ mag (Tables 1 and 3), which is too faint to be stars at that distance.

Fourth, the NIR to X-ray flux ratio of group A sources ($F_{K_S}/F_X \sim 10^1\text{--}10^4$) is consistent with typical values of the X-ray to the bolometric luminosity ratio of $L_X/L_{\text{bol}} \sim 10^{-5}\text{--}10^{-2}$ (Preibisch et al. 2005) of PMS sources. Assuming an age of 2 Myr (the same with IC 348; Muench et al. 2003), and using $L_{K_S}/L_{\text{bol}} \sim 4\text{--}9 \times 10^{-2}$ ($0.08\text{--}1.4 M_\odot$ PMS source; Baraffe et al. 1998; D'Antona & Mazzitelli 1994) and $L_{K_S}/L_X \sim F_{K_S}/F_X \sim$




Fig. 5.— Color-magnitude diagram of X-ray sources with NIR counterparts (*filled squares*) and FLAMINGOS sources (*open circles*). Sources with J -, H -, and K_S -band detections with magnitude uncertainty of < 0.05 mag are plotted. Labels are given to the X-ray sources with NIR counterparts. The 1, 2, and 3 Myr isochrone curves (*dotted*, *solid*, and *dashed curves*) are from Baraffe et al. (1998). The reddening lines for $1.4 M_\odot$, $0.2 M_\odot$, and $0.08 M_\odot$ are given with dashed lines assuming an age of 2 Myr.

5×10^2 (the median value of group A sources), we obtain the value of $L_X/L_{\text{bol}} \sim 0.8\text{--}1.8 \times 10^{-4}$ for group A sources. A similar L_X/L_{bol} value is obtained if we assume that these sources are main sequence stars (Tokunaga 2000; Drilling & Landolt 2000), which is higher than their typical values ($10^{-7}\text{--}10^{-5}$; Preibisch et al. 2005). We thus consider that most of the group A sources are YSOs in the cloud and not main sequence stars in the line of sight. The distinction between YSO and main sequence natures should be confirmed with follow-up spectroscopic studies. Group B sources, on the other hand, have F_{K_S}/F_X values of $10^{-1}\text{--}10^1$, which is consistent with the values seen in an X-ray-selected sample of AGNs (Watanabe et al. 2004).

Figure 6 shows a color-color diagram of NIR-identified X-ray (*squares*) and FLAMINGOS (*circles*) sources. All NIR-identified X-ray sources in group A (Nos. 4, 6, 7, 14, 15, 23, 30, 42, and 48) have NIR colors consistent with being reddened weak-line T Tauri or main sequence stars. None of them shows apparent NIR excess emission, which does not exclude the possibility that they are classical T Tauri stars or protostars, as K_S -band observations are not completely sensitive to excess emission from circumstellar disks (e.g., Haisch et al. 2001). The only source with significant J -, H -, and K_S -band detections in group B (No. 16) has an inconsistently red color for reddened stars of $m_J - m_{K_S} = 2.0$ mag, which is commonly seen among 2MASS red AGNs (Cutri et al. 2001). The density of such sources is $\sim 0.5 \text{ arcdeg}^{-2}$ at $K_S < 14.5$ mag (Cutri et al. 2001), making the probability $\sim 5\%$ to have a 2MASS red AGN in an ACIS field.

In total, twelve X-ray sources (Nos. 4, 6, 7, 14, 15, 23, 30, 34, 42, 48, 68, and 69) have X-ray and NIR features consistent of being low-mass YSOs in this cloud. Two of them are known sources; No. 7 is IRS 1 and No. 42 is an A5 type star (Roeser & Bastian 1988) that consists of a double (Worley & Douglass 1997). Main sequence A type stars are not expected to have X-ray emission due to the lack of any plasma creation mechanisms (Schmitt et al. 1993). The X-ray emission in No. 42 may come from a low-mass companion, which mimics a F_{K_S}/F_X value consistent with a PMS source (Fig. 5). The remaining ten sources are candidates of new YSOs in L1448. The brightness histogram of the rest of the X-ray sources is consistent with

a log N -log S relation of AGNs (Moretti et al. 2003).




Fig. 6.— Color-color diagram of X-ray sources with NIR counterparts (*filled squares*) and FLAMINGOS sources (*open circles*). Sources with J -, H -, and K_S -band detections with magnitude uncertainty of < 0.1 mag are plotted. FLAMINGOS sources with NIR excess emission are shown with larger symbols. The reddening lines are shown (*dashed lines*) from the intrinsic colors of dwarfs and giants (*thick solid curves*; Tokunaga 2000) and the classical T Tauri star locus (*thick solid lines*; Meyer et al. 1997).

4.2. X-ray and NIR Emission from Embedded Sources

4.2.1. Infrared Excess Sources

From Figure 6, ten FLAMINGOS sources (*large open circles*) are found to have NIR excess emission and thus are candidates for new embedded sources. Daggers are given for these sources in Table 3. Among them, FLAMINGOS Nos. 23 and 149 are promising YSO candidates with estimated magnitude uncertainty of $\lesssim 0.05$ mag.

4.2.2. IRS 1

Both X-ray and NIR emission was found from IRS 1 (ACIS No. 7 in Table 1 and FLAMINGOS

No. 2 in Table 3 or 2MASS 03250943+3046215). The X-ray and NIR properties are consistent with the nature of this source as a classical T Tauri star or a class I protostar; (1) it shows X-ray flux variation commonly seen in X-ray emission of stellar magnetic activity origin, (2) its ME and F_{Ks}/F_X values indicate this source to be a YSO (Fig. 4), and (3) a nebulous structure is seen in the FLAMINGOS K_S -band image (Fig. 1b). The NIR magnitudes of this source ($m_J \sim 12.5$ mag, $m_H \sim 11.0$ mag, and $m_{Ks} \sim 10.1$ mag) correspond to an object with the mass of $0.5\text{--}1.0 M_\odot$ at 300 pc obscured by $A_V \sim 7$ mag.

4.2.3. IRS 3(A)

No emission is seen at the position of IRS 3(A) in our NIR images. The 5σ level of the local sky noise corresponds to ~ 18.8 mag, ~ 17.7 mag, and ~ 17.5 mag in our J -, H -, and K_S -band images. The SED slope defined as $\alpha = (d \log \lambda F_\lambda) / (d \log \lambda)$ between $2.2 \mu\text{m}$ (K_S band) and $10 \mu\text{m}$ (N band) is commonly used to classify YSOs (Wilking et al. 1989), using the fact that the $2\text{--}10 \mu\text{m}$ SEDs at a younger evolutionally stage are more dominated by emission from circumstellar material. The lower limit of α of IRS 3(A) is $\gtrsim 3.2$ from the non-detection in the NIR bands and a MIR detection in the N band (Ciardi et al. 2003). Here, we calculated the α value as the index between H and N bands, as the K_S -band image is noisy at the position due to the emission arising from the outflow of IRS 3(A) itself. The value thus is a lower limit also in the sense that H -band emission is subject to contamination from reflected photospheric emission.

Greene et al. (1994) gave a qualitative classification of YSOs using α , where $\alpha < -1.6$ for class III, $-1.6 < \alpha < -0.3$ for class II, and $\alpha > -0.3$ for class I sources. Almost all known class I protostars have $\alpha \lesssim 2$ with the only exception of WL 22 ($\alpha = 3$) in ρ Ophiuchi dark cloud (Wilking et al. 1989). Sources with a steeper index than ~ 3 have been rarely known, as they are quite difficult to detect in K band images that traditionally pilot protostar searches. However, recent progress of sensitive MIR imaging both by ground- and space-based telescopes is changing the situation by reporting several protostars with very steep spectra with indices of $\alpha > 3$ (Cep E by Noriega-Crespo et al. 2004 and source X_E

in R CrA by Hamaguchi et al. 2005 and private communications with K. Nedachi). The index of $\gtrsim 3.2$ of IRS 3(A) places this source in the class of very steep spectrum protostars. More sources in this emerging class are expected to be discovered with *Spitzer* observations of near-by star-forming regions. Some of them are class 0 protostars (e.g., Cep E mm; Noriega-Crespo et al. 2004) and others are K -band-unidentified class I protostars, or in between.

An unusual concentration of three X-ray photons within a $\sim 0.5''$ radius ($\sim 90\%$ encircled energy radius at $\sim 0.4'$ off-axis angle) can be found at the position of IRS 3(A). No such concentration is seen within $1'$ of the on-axis position. Although it was not recognized as a source in the initial X-ray source search using the wavelet technique, its PNS value of 5×10^{-4} indicates the existence of an X-ray source of $F_X \approx 5 \times 10^{-16} \text{ ergs s}^{-1} \text{ cm}^{-2}$. The facts that the position of the concentration is consistent with that of IRS 3(A), that the energy of all photons is concentrated in the hard (2–5 keV) band, and that the arrival times of all photons fall in a limited time range of ~ 15 ks in the ~ 68 ks observation, which is indicative of a flare, may further strengthen the claim. There is no indication of temporal non-uniformity of background counts. A similar weak but plausible X-ray emission was found in an embedded source (FIR4) in NGC 2024 by *Chandra* (Skinner et al. 2003).

Given the capability of *Chandra* and MIR telescopes (*Spitzer* and ground-based facilities), it is now becoming possible to investigate X-ray emission from protostars with a very steep spectrum ($\alpha > 3$) even without K_S -band identifications. If IRS 3(A) is indeed an X-ray source, it has the steepest spectral index among YSOs with X-ray emission. Although it is not yet certain whether a steeper α value indicates a younger age of protostars, X-ray observations for steep spectrum sources are a key to answer how early X-ray emission starts and how intense it is at the earliest stage of star formation. These are important questions to understand star formation because the coupling between gas and magnetic fields can be largely affected by X-ray ionization (Feigelson & Montmerle 1999).

4.2.4. Class 0 Sources

No significant X-ray or NIR emission was found from the four known class 0 sources; L1448 mm, IRS 2, IRS 3(B), and IRS 3(C). The upper limit ($PNS = 1 \times 10^{-3}$) of the X-ray flux in the 0.5–8.0 keV is $\sim 0.5\text{--}1.0 \times 10^{-15}$ ergs s cm $^{-2}$, whereas the upper limits (5σ) in the NIR regime are ~ 18.8 mag (J), ~ 17.7 mag (H), and ~ 17.5 mag (K_S), except for L1448 mm and IRS 3(B) in the K_S -band, which are contaminated by local diffuse emission.

The lack of X-ray detection from class 0 sources is widely reported. Froebrich (2005) compiled 28 well documented bona-fide class 0 sources. Sixteen of them were observed by *Chandra* and nine in references are reported to have no X-ray detection (L1448 NW and C; this work, NGC 1333 I2, I4A, and I4B; Getman et al. 2002, OMC 3 MMS6; Tsujimoto et al. 2002, HH 24 MMS and HH 25 MMS; Simon et al. 2004, and VLA 1623; Imanishi 2003). An additional six sources are reported with no detection in *XMM-Newton* references (IRAS 03256+3055; Preibisch & Zinnecker 2002, HH 211 MM; Preibisch & Zinnecker 2004, Serpens S68N, SMM2, SMM3, and SMM4; Preibisch 2004). Considering that these sources are deeply embedded, the lack of X-ray detection does not necessarily indicate that they do not have any X-ray-emitting activities, although they are still difficult for X-ray as well as MIR investigations with current technologies.

The authors express gratitude for Kentaro Motohara for his help in the observation at Kitt Peak, Pat Broos for his advice in X-ray data reduction using ACIS Extract and proofreading, and Eric D. Feigelson for discussion. We acknowledge financial supported by the Japan Society for the Promotion of Science (M. T.), a Grant-in-Aid for Scientific Research of the Ministry of Education, Culture, Sports, Science and Technology (No. 15740120; Y. T.), a Chuo University Grant for Special Research (Y. T.), the Saneyoshi Scholarship Foundation (Y. T.), and the *Chandra* guest observer grant (SAO G04–5009X). FLAMINGOS was designed and constructed by the IR instrumentation group (PI: R. Elston) at the University of Florida, Department of Astronomy, with support from NSF grant AST97-31180 and Kitt Peak Na-

tional Observatory. IRAF is distributed by the National Optical Astronomy Observatories, which are operated by the Association of Universities for Research in Astronomy, Inc., under cooperative agreement with the National Science Foundation. This publication makes use of data products from the Two Micron All Sky Survey, which is a joint project of the University of Massachusetts and the Infrared Processing and Analysis Center/California Institute of Technology, funded by the National Aeronautics and Space Administration and the National Science Foundation.

Facilities: CXO(ACIS), KPNO:4m(FLAMINGOS)

REFERENCES

- Anglada, G., Rodriguez, L. F., Torrelles, J. M., Estalella, R., Ho, P. T. P., Canto, J., Lopez, R., & Verdes-Montenegro, L. 1989, *ApJ*, 341, 208
- Anglada, G., & Rodríguez, L. F. 2002, *Revista Mexicana de Astronomia y Astrofisica*, 38, 13
- Bachiller, R., & Cernicharo, J. 1986, *A&A*, 168, 262
- Bachiller, R., Martin-Pintado, J., Tafalla, M., Cernicharo, J., & Lazareff, B. 1990, *A&A*, 231, 174
- Bachiller, R., Andre, P., & Cabrit, S. 1991, *A&A*, 241, L43
- Bally, J., Lada, E. A., & Lane, A. P. 1993, *ApJ*, 418, 322
- Bally, J., Devine, D., Alten, V., & Sutherland, R. S. 1997, *ApJ*, 478, 603
- Baraffe, I., Chabrier, G., Allard, F., & Hauschildt, P. H. 1998, *A&A*, 337, 403
- Barsony, M., Ward-Thompson, D., André, P., & O’Linger, J. 1998, *ApJ*, 509, 733
- Bertin, E. & Arnouts, S. 1996, *A&AS*, 117, 393
- Ciardi, D. R., Telesco, C. M., Williams, J. P., Fisher, R. S., Packham, C., Piña, R., & Radomski, J. 2003, *ApJ*, 585, 392
- Cohen, M., & Kuhi, L. V. 1979, *ApJS*, 41, 743
- Curiel, S., Raymond, J. C., Moran, J. M., Rodriguez, L. F., & Canto, J. 1990, *ApJ*, 365, L85

- Cutri, R. M., et al. 2001, *Astronomical Society of the Pacific Conference Series*, 232, 78
- D’Antona, F., & Mazzitelli, I. 1994, *ApJS*, 90, 467
- Davis, C. J., & Smith, M. D. 1995, *ApJ*, 443, L41
- de Grijp, M. H. K., Lub, J., & Miley, G. K. 1987, *A&AS*, 70, 95
- Drilling, J. S. & Landolt, A. U. 2000, in *Allen’s Astrophysical Quantities*, ed. A. N. Cox (New York: Springer-Verlag), 381
- Eisloffel, J. 2000, *A&A*, 354, 236
- Elston, R., Raines, S. N., Hanna, K. T., Hon, D. B., Julian, J., Horrobin, M., Harmer, C. F. W., & Epps, H. W. 2003, *Proc. SPIE*, 4841, 1611
- Feigelson, E. D., & Montmerle, T. 1999, *ARA&A*, 37, 363
- Froebrich, D. 2005, *ApJS*, 156, 169
- Garmire, G. P., Bautz, M. W., Ford, P. G., Nousek, J. A., & Ricker, G. R. 2003, *Proc. SPIE*, 4851, 28
- Gehrels, N. 1986, *ApJ*, 303, 336
- Getman, K. V., Feigelson, E. D., Townsley, L., Bally, J., Lada, C. J., & Reipurth, B. 2002, *ApJ*, 575, 354
- Getman, K. V., et al. 2005, *ApJS*, in press
- Greene, T. P., Wilking, B. A., Andre, P., Young, E. T., & Lada, C. J. 1994, *ApJ*, 434, 614
- Haisch, K. E., Lada, E. A., Piña, R. K., Telesco, C. M., & Lada, C. J. 2001, *AJ*, 121, 1512
- Hamaguchi, K., Corcoran, M. F., Petre, R., White, N. E., Stelzer, B., Nedachi, K., Kobayashi, N., & Tokunaga, A. T. 2005, *ApJ*, 623, 291
- Imanishi, K. 2003, Ph.D. thesis, Kyoto University
- Krautter, J., et al. 1999, *A&A*, 350, 743
- Lynds, B. T. 1962, *ApJS*, 7, 1
- Maccacaro, T., Gioia, I. M., Wolter, A., Zamorani, G., & Stocke, J. T. 1988, *ApJ*, 326, 680
- Meyer, M. R., Calvet, N., & Hillenbrand, L. A. 1997, *AJ*, 114, 288
- Monet, D. G., et al. 2003, *AJ*, 125, 984
- Moretti, A., Campana, S., Lazzati, D., & Tagliaferri, G. 2003, *ApJ*, 588, 696
- Muench, A. A., et al. 2003, *AJ*, 125, 2029
- Noriega-Crespo, A., Moro-Martin, A., Carey, S., Morris, P. W., Padgett, D. L., Latter, W. B., & Muzerolle, J. 2004, *ApJS*, 154, 402
- O’Linger, J., Wolf-Chase, G., Barsony, M., & Ward-Thompson, D. 1999, *ApJ*, 515, 696
- Preibisch, T., Zinnecker, H., & Herbig, G. H. 1996, *A&A*, 310, 456
- Preibisch, T., & Zinnecker, H. 2001, *AJ*, 122, 866
- Preibisch, T., & Zinnecker, H. 2002, *AJ*, 123, 1613
- Preibisch, T. 2003, *A&A*, 401, 543
- Preibisch, T. 2004, *A&A*, 428, 569
- Preibisch, T., & Zinnecker, H. 2004, *A&A*, 422, 1001
- Preibisch, T., et al. 2005, *ApJ*, submitted
- Roeser, S., & Bastian, U. 1988, *A&AS*, 74, 449
- Scargle, J. D. 1998, *ApJ*, 504, 405
- Schmitt, J. H. M. M., Zinnecker, H., Cruddace, R., & Harnden, F. R. 1993, *ApJ*, 402, L13
- Simon, T., Andrews, S. M., Rayner, J. T., & Drake, S. A. 2004, *ApJ*, 611, 940
- Skinner, S., Gagné, M., & Belzer, E. 2003, *ApJ*, 598, 375
- Skrutskie, M. F., et al. 1997, *ASSL Vol. 210: The Impact of Large Scale Near-IR Sky Surveys*, 25
- Smith, R. K., Brickhouse, N. S., Liedahl, D. A., & Raymond, J. C. 2001, *ApJ*, 556, L91
- Tokunaga, A. T. 2000, in *Allen’s Astrophysical Quantities*, ed. A. N. Cox (New York: Springer-Verlag), 143
- Tsujimoto, M., Koyama, K., Tsuboi, Y., Goto, M., & Kobayashi, N. 2002, *ApJ*, 566, 974

- Watanabe, C., Ohta, K., Akiyama, M., & Ueda, Y. 2004, *ApJ*, 610, 128
- Weisskopf, M. C., Brinkman, B., Canizares, C., Garmire, G., Murray, S., & Van Speybroeck, L. P. 2002, *PASP*, 114, 1
- Wilking, B. A., Lada, C. J., & Young, E. T. 1989, *ApJ*, 340, 823
- Worley, C. E., & Douglass, G. G. 1997, *A&AS*, 125, 523
- Yamauchi, S., Tsuboi, Y., & Koyama, K. 2001, *PASJ*, 53, 885